

X-ray Crystallography from a Quantum Mechanical Perspective

Frank Rioux
Department of Chemistry
St. John's University/College of St. Benedict


According to Richard Feynman¹ the double-slit experiment is the paradigm for all of quantum mechanics because it is a simple manifestation of the superposition principle, which is quantum theory's "only mystery." In an interesting quantum mechanical analysis of Young's double-slit experiment, Marcella² writes the coordinate wave function of the particle interacting with the slit-screen as a linear superposition of being at both slits,

$$|\Psi\rangle = \frac{1}{\sqrt{2}} [|x_1\rangle + |x_2\rangle] \quad (1)$$


where x_1 and x_2 are the positions of two infinitesimally thin slits. The diffraction pattern at the detection screen is determined by the momentum distribution at the slit screen. The momentum wave function is obtained by projecting equation (1) into momentum space,

$$\langle p_x | \Psi \rangle = \frac{1}{\sqrt{2}} [\langle p_x | x_1 \rangle + \langle p_x | x_2 \rangle] = \frac{1}{2\sqrt{\pi\hbar}} \left[\exp\left(-\frac{ip_x x_1}{\hbar}\right) + \exp\left(-\frac{ip_x x_2}{\hbar}\right) \right] \quad (2)$$

where the $\langle p_x | x_i \rangle$ are the position eigenfunctions in the momentum representation. The square of the absolute magnitude of this function is shown in Figure 1.



Marcella improves this model by giving the slits a finite width, yielding the more realistic diffraction pattern shown in Figure 2.



The uncertainty principle is clearly illustrated in both of these simple models. The particle has been sharply localized in coordinate space by the slits, therefore its momentum must be delocalized as shown in Figures 1 and 2. In addition, because the particle has been localized at two positions interference effects are prominent in its momentum distribution.

Like the double-slit phenomenon, x-ray diffraction by crystals is also based on the linear superposition principle. For the sake of simplicity attention will be restricted to two-dimensional crystals. Also, following Marcella's first model, the atoms or ions occupying the lattice sites are considered to be point scatterers. Under these assumptions, the coordinate-space wave function of a photon illuminating a crystal is a superposition of the atomic scattering positions.

$$|\Psi\rangle = \frac{1}{\sqrt{N}} \sum_{i=1}^N |x_i, y_i\rangle \quad (3)$$


As we wish to know the diffraction pattern that will be found at the detector, we express this superposition in the momentum representation as follows.

$$\langle p|\Psi\rangle = \frac{1}{\sqrt{N}} \sum_{i=1}^N \langle p|x_i, y_i\rangle = \frac{1}{\sqrt{N}} \sum_{i=1}^N \langle p_x|x_i\rangle \langle p_y|y_i\rangle \quad (4)$$

Employing the expression for the position wave function in the momentum representation we can convert equation (4) to

$$\langle p|\Psi\rangle = \frac{1}{\sqrt{N}(2\pi\hbar)} \sum_{i=1}^N f_i \exp\left[-\frac{i}{\hbar}(p_x x_i + p_y y_i)\right] \quad (5)$$

where the f_i are the atomic scattering factors. A face-centered “squaric” arrangement of five atomic point scatterers yields the following diffraction pattern, $|\langle p|\Psi\rangle|^2$.



P

The Mathcad document used to generate this diffraction pattern is shown in the Appendix.

Literature cited:

1. R. P. Feynman, R. B. Leighton, and M. Sands, *The Feynman Lectures on Physics*, Vol. 3; Addison-Wesley: Reading, 1965.
2. T. V. Marcella, "Quantum interference with slits," *European Journal of Physics* **23**, 615-621 (2002).

Appendix

Number of atoms: $A := 5$ Atomic positions and relative scattering factors:

$$x_1 := 0 \quad y_1 := 0 \quad f_1 := \frac{1}{4} \quad x_2 := 0 \quad y_2 := 1 \quad f_2 := \frac{1}{4}$$

$$x_3 := 1 \quad y_3 := 0 \quad f_3 := \frac{1}{4} \quad x_4 := 1 \quad y_4 := 1 \quad f_4 := \frac{1}{4}$$

$$x_5 := \frac{1}{2} \quad y_5 := \frac{1}{2} \quad f_5 := \frac{1}{2}$$

$$\Delta := 10 \quad N := 200 \quad j := 0..N \quad p_{x_j} := -\Delta + \frac{2 \cdot \Delta \cdot j}{N} \quad k := 0..N \quad p_{y_k} := -\Delta + \frac{2 \cdot \Delta \cdot k}{N}$$

Fourier transform of position wave function into the momentum representation:

$$\Psi(p_x, p_y) := \sum_{m=1}^A f_m \cdot \exp(-i \cdot p_x \cdot x_m - i \cdot p_y \cdot y_m) \quad P_{j,k} := \left(\left| \Psi(p_{x_j}, p_{y_k}) \right| \right)^2$$